

SMALL-BODIED FISHES MONITORING IN THE SAN JUAN RIVER

FY 2022 SCOPE OF WORK

TO

BUREAU OF RECLAMATION

FROM

New Mexico Department of Game and Fish
Adam L. Barkalow and Matthew P. Zeigler
One Wildlife Way, P.O. Box 25112
Santa Fe, New Mexico 87504
505-476-8104
adam.barkalow@state.nm.us
matthew.zeigler@state.nm.us

SUBMITTED

26 March, 2021

FOR

Title of Agreement: Small-bodied Fishes Monitoring on the San Juan River
Agreement Number: R19AC00022

REPORTING DATES:

10/01/2021 through 9/30/2022

This scope of work (SOW) is for Small-bodied Fishes (SBF) Monitoring within the San Juan River for FY2022. This project aims to provide the highest quality data and quantitative assessment on the effects of management actions on the survival of post-larval early life stages of native and nonnative fishes and their recruitment into subsequent life stages and use this information to recommend appropriate modifications to recovery strategies for Colorado Pikeminnow and Razorback Sucker in the San Juan River.

NEED

The San Juan River Basin Recovery Implementation Program (SJRIP) Long-Range Plan specifies that monitoring and evaluation of fish is a necessary element for assessing the recovery of federally endangered Colorado Pikeminnow *Ptychocheilus lucius* and Razorback Sucker *Xyrauchen texanus* in the San Juan River (Element 4; SJRIP 2016). Task 4.1.2.2 of the SJRIP's Long-Range Plan specifies the need for juvenile and small-bodied fishes (SBF) monitoring to locate areas and habitats used for rearing and to determine if young fish are surviving and recruiting into adult populations (SJRIP 2016). Data collected during annual SBF monitoring is used to assess recovery of Colorado Pikeminnow and Razorback Sucker, as well as evaluate the influences of SJRIP management actions on the river's fish community as a whole (Gido and Propost 2012; Franssen et al. 2015; Zeigler and Ruhl 2017).

GOAL

The goal of SBF monitoring is to quantitatively assess the effects of management actions on survival of post-larval early life stages of native and nonnative fishes and their recruitment into subsequent life stages and use this information to recommend appropriate modifications to recovery strategies for Colorado Pikeminnow and Razorback Sucker in the San Juan River (SJRIP 2012).

MONITORING OBJECTIVES

The specific objectives for SBF monitoring include:

1. Annually document occurrence and density of native and nonnative age-0/small-bodied fishes in the San Juan River.
2. Document mesohabitat use by age-0 Colorado Pikeminnow, Razorback Sucker, and Roundtail Chub, as well as other native and nonnative fishes in the primary channel, secondary channels, and backwaters.
3. Obtain data that will aid in the evaluation of the responses of native and nonnative fishes to different flow regimes and other management actions.
4. Track trends in native and nonnative fish populations.
5. Characterize patterns of mesohabitat use by native and nonnative small-bodied fishes.

STUDY AREA

The spatial extent of small-bodied fishes monitoring has varied since 2003 (Figure 1). Until 2011, sampling occurred every year from River Mile (RM) 180.6 (Animas River confluence) downstream to RM 2.9 (Clay Hills Crossing, UT) (Figure 2). Sampling below RM 76.4 (Sand Island, UT) occurred once every five years after 2010, primarily because the fish fauna in geomorphic Reaches 1 and 2 was mostly (~80%) nonnative fishes and had shown little change since the initiation of SBF monitoring (Zeigler and Ruhl 2016). Sampling was extended upstream to RM 196.0 (Bloomfield, NM) in 2012. Beginning in 2017, sampling area was determined using a flexible schematic where different sections of the river were sampled based on the number of wild age-0 Colorado Pikeminnow or Razorback Sucker captured during sampling from RM 147.8 (Shiprock, NM) downstream to RM 52.7 (Mexican Hat, UT). Due to the unique detections of wild age-0 Razorback Sucker and increasingly consistent capture of Colorado Pikeminnow, especially in downstream reaches (Figure 3), a downstream expansion of SBF monitoring (Reaches 1 and 2) was proposed in FY2020 and FY2021 SOWs. (Zeigler and Wick 2019; Barkalow et al. 2020). Small-bodied fishes Monitoring in 2020 was approved to sample from RM 147.8 to RM 2.9 but sampling was restricted to between RM 76.4 and RM 2.9 due to COVID-19 related closures (Figure 1). Small-bodied

fishes monitoring detected wild age-0 Colorado Pikeminnow in Reach 1 for the first time ever in 2020, demonstrating that sampling in the lower reaches may lends insight into habitat use and distribution of wild spawned Colorado Pikeminnow in the San Juan River (Barkalow and Zeigler in preparation).

Sampling in 2022 will occur from RM 180.6 downstream to RM 2.9 (Figure 2). Recent captures of wild age-0 endangered fishes in downstream reaches (1– 4) indicates that sampling Reaches 1– 6 consistently may be more advantageous to meeting the SBF monitoring objectives and project goals (Zeigler and Ruhl 2017; Barkalow and Zeigler 2021 in preparation). Additionally, endangered fish have not been captured upstream of RM 180.6 during SBF monitoring. Sampling from RM 180.6 downstream to RM 3.0 will ensure the continuity of the long-term data set for Reaches 3 – 6 and will align SBF monitoring with observed patterns of larval endangered fish densities while maintaining sampling within areas that have produced the highest wild age-0 Colorado Pikeminnow captures (Figures 5 and 11 in Farrington et al. 2020). The proposed section of river to be sampled will cover both the entire reach sampled during the Demographic Monitoring project and be congruent with the proposed post-2023 sampling regime. The spatial extent of area sampled could be expanded to include other areas (e.g., Lake Powell or Animas River) on a yearly basis based on new evidence of increased endangered species presence in the areas above or below the sampling area.





Figure 1. Spatial extent of sampling during small-bodied fishes monitoring on the San Juan River since 2003. The spatial extent of sampling in 2021 and 2022 are as proposed. Note that river miles begin at the inflow of Lake Powell in Utah (River Mile 0) and end at Navajo Dam in New Mexico (River Mile 224).



1
2 Figure 2. The San Juan River in New Mexico, Colorado, and Utah with river miles and geomorphic reaches indicated. The section between the red arrows will be sampled during
3 2022 small-bodied fishes monitoring (Animas River confluence downstream to Clay Hills Crossing, UT). Inset indicates location of San Juan River in Colorado, New Mexico, and
4 Utah.




Figure 3. The expected catch-per-unit-effort (ECPUE, fish/10 m²) of wild age-0 Colorado Pikeminnow in Reaches 1 – 7 in 2016, 2017, and 2019 overlaid on the extent in percent of total Reach length sampled in the same years. Note that Reach 1 was not sampled in those years. Circles represent mean ECPUE estimates and bars represent 95% confidence interval; gray bars indicate percent of extent sampled within each reach.

METHODS

Small-bodied fishes monitoring is designed to sample habitats which have the greatest likelihood of supporting age-0 large-bodied fishes (e.g., Bluehead Sucker *Catostomus discobolus* and Flannelmouth Sucker *Catostomus latipinnis*) and all age classes of small-bodied fishes (e.g., Red Shiner *Cyprinella lutrensis* and Speckled Dace *Rhinichthys osculus*). Sampling will occur at designated 3-mile intervals in the primary channel, and at all secondary (i.e., less than 20% of total flow) and all zero velocity channels (i.e., backwaters and embayments; ≥ 30 m²) when encountered (SJRIP 2012). Sample reaches will be approximately 200 m long (measured along the shoreline) at primary channel sample sites and, depending upon the extent of surface water, 100 – 200 m long at secondary and zero velocity channel sample sites.

In the fall of 2012, six secondary channels were modified during the Phase I River Ecosystem Restoration Initiative (RERI) habitat restoration efforts through excavation of sediment and removal of nonnative plants. These channels are located at RM 132.2, 132.0, 130.7A, 130.7B, 128.6, and 127.2. An additional channel located at RM 136.5 was restored during Phase II habitat restoration efforts in 2014. These restoration sites will be visited during annual SBF monitoring and sampled if flowing following the protocols described below.

River mile, geographic coordinates (UTM NAD83), and water quality parameters (dissolved oxygen, conductivity, and temperature) will be recorded at each sampling site. All mesohabitats (e.g., riffle, run, pool) present within a site (except large zero velocity channels) will be sampled in rough proportion to their availability using a 3.0 x 1.8 m (3.0 mm heavy duty Delta untreated mesh) drag seine. Seine hauls will be made in at least eight different mesohabitats at each site; however, if habitat heterogeneity is low at a site, as few as five seine hauls will be made. At least two seine hauls, one across the mouth and one parallel to its long axis will be made at each large zero velocity channel unless the mouth is too narrow, in which case only one seine haul, parallel to its long axis, will be made.

All captured fishes will be identified to species, enumerated, and up to 25 age-0 and age-1+ fish will be measured for total length (mm TL). All fish that are not individually measured will be recorded by age class (age-0 or age-1+) and minimum and maximum total length (mm TL) will be recorded for those fishes. Any captured endangered or rare species (i.e., Colorado Pikeminnow, Razorback Sucker, and Roundtail Chub *Gila robusta*) will be measured for both standard and total length (mm SL, TL), weighed (g). Endangered or rare fish ≥ 130 mm will be interrogated for a passive integrated transponder (PIT) tag and if one is not detected a 12 mm PIT tag will be injected into the intramuscular tissue below the pelvic girdle. All native fishes will be released and nonnative fishes removed from the river. Fishes too small to easily identify in the field will be fixed in 10% solution of formalin and returned to the laboratory for identification. Tissue samples (fin clip) will be collected from any juvenile (≤ 300 mm TL) untagged Razorback Sucker captured during SBF monitoring for genetic analysis. Fin clips will be stored in individually labeled centrifuge tubes that contain 100% ethanol.

Following each seine haul, the sampled width and length of each mesohabitat will be measured to the nearest 0.1 m and recorded. The depth and dominant substrate at five generalized locations, and any cover (e.g., boulders, debris piles, large woody debris) associated with the mesohabitat will also be recorded. Retained specimens and tissue samples will be identified and measured (TL and SL) in the laboratory to the nearest 0.1 mm and accessioned to the University of New Mexico Museum of Southwest Biology (MSB), Division of Fishes.

After data collection, all original field notes will be checked for errors and missing data. Data will be entered into Excel spreadsheets with a similar template as a project specific database. All entered data will be cross-checked with the original field notes by a different biologist. Data from the Excel spreadsheets will be imported into the database. Specific conditions for each data field in the database prevent the entry of incorrect data and typographical errors. Database queries will be used to identify and rectify any additional errors. Following the submission of the final report original field notes will be deposited to MSB Division of Fishes.

DATA ANALYSIS AND REPORTING

Analyses will be based on density (i.e., catch-per-unit-effort, CPUE) of individual species, calculated by seine haul, as the number of fish captured per square meter sampled (width x length). To account for the significant number of zeros and highly skewed data, density for each species will be analyzed using a Delta-GLM approach which combines two separate components: (1) a logistic model estimating the probability of presence ($CPUE_{0/1}$) fitted using a GLM with a binomial distribution and logit link, and (2) a model for CPUE only when the species is present ($CPUE^+$) fitted using a GLM with a lognormal distribution (Fletcher et al. 2005; Acou et al. 2011; Vasconcelos et al. 2013). The predicted density, $E(CPUE)$, is then obtained by (3) multiplying the response variables predicted by the binomial and lognormal models for each individual seine haul. This approach models the two aspects of the data

(i.e., presence/absence and positive density) separately, allowing for evaluation of how covariates influence the two separate processes. Furthermore, the approach is much simpler and easier to interpret than other methods such as mixture models (Fletcher et al. 2005). Calculation of density will be limited to those species which have greater than 3% of all seine hauls (2003 – 2021) with at least one capture.

Several models using a combination of covariates will be used in both the logistic and positive lognormal model. Both abiotic and biotic covariates will be investigated for their use in each model (Table 1). The negative log-likelihood from both models will be combined to calculate Akaike’s Information Criterion with a correction for finite sample sizes (AIC_c). The combined model with the lowest AIC_c will then be used to model the final binomial and lognormal models for each species. Residual plots will be examined to ensure that the final positive lognormal model meets the assumptions of normally distributed and equal variance residuals.

Table 1. Name and description of abiotic and biotic covariates which may be used in both the logistic and lognormal models to predicted expected density (E(CPUE)).

Covariate Name	Description
sampYear	Year in which the sample was taken.
Reach	Geomorphic reach in which the sampled was taken.
RiverMile	The river mile where the sample was taken.
ChannelType	The channel type in which the sampled was taken.
Mesohabitat	The mesohabitat in which the sample was taken.
sampDis	Discharge at time the sample was taken.
NNC_1_Den	The density of nonnative competitors in the Reach where the sample was taken. Calculated as the total density of Fathead Minnows, Red Shiners, and Western Mosquitofish captured during annual small-bodied fishes monitoring.
NNC_2_Den	The density of nonnative competitors in the Reach where the sample was taken. Calculated as the total density of Channel Catfish, Fathead Minnows, Red Shiners, and Western Mosquitofish captured during annual small-bodied fishes monitoring.
NNC_3_Den	The density of nonnative competitors in the Reach where the sample was taken. Calculated as the total density of Channel Catfish captured during annual small-bodied fishes monitoring.

Final binomial, log-normal, and Delta-GLM models will be assessed for goodness-of-fit and predictive capability. The predictive accuracy for the binomial model will be tested using the Area Under the Curve (AUC) of the Receiver Operating Characteristic (ROC). The ROC analysis involves plotting the proportion of known presences predicted against the proportion of known absences predicted (Peterson et al. 2008). The values of the AUC of the ROC curve range from 0.5 to 1.0 with 0.5 indicating no fit and a 1.0 a perfect fit (Fielding and Bell 1997). For the positive lognormal and Delta-GLM models, a linear regression between observed (x-axis) and predicted (y-axis) CPUE will be used to test predictive ability. The coefficient of determination (R²) of this relationship shows the proportion of the linear variation in y (predicted values) explained by the variation in x (observed values), the intercept of this linear regression model describes bias, and the slope describes consistency.

Data collected from the six RERI and the Phase II secondary channels will be reported for each year since these channels were restored. Information for the RERI and Phase II secondary channels will

include if the channel was sampled, reasoning for why it was not sampled, and number of endangered, native, and nonnative species captured.

An annual report will provide a summation of data obtained in FY2022, a synthesis of data across years to document/assess species populations' trends, and a summary of mesohabitat associations. Separate data summaries and analyses will also be conducted for any wild age-0 CPM and RBS, if needed. All data collected will be recorded on electronic spreadsheets and provided to USFWS Program Office by the principal investigator, along with the annual final report, by 30 June 2023.

Budget Justification

The FY2022 SOW budget is a slightly different cost from FY2021 due to a recent increase in New Mexico state employee per diem rates. Additionally, the negotiated IDC rate is slightly lower than 2021. There were no other modifications to the budget.

REFERENCES

- Acou, A., E. Rivot, J. A. Van Gils, A. Legault, F. Ysnel, and E. Feunteun. 2011. Habitat carrying capacity is reached for the European eel in a small coastal catchment: evidence and implications for managing eel stocks. *Freshwater Biology* 56:952-968.
- Barkalow, A.L., M.P. Zeigler, J.M. Wick. 2020. Small-bodied fishes monitoring in the San Juan River: 2019. Final Report submitted to the San Juan River Basin Recovery Implementation Program. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico
- Barkalow, A.L. and M.P. Zeigler. 2021 in preparation. Small-bodied fishes monitoring in the San Juan River: 2020. Draft Report in preparation to be submitted to the San Juan River Basin Recovery Implementation Program. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- Farrington, M. A., R. K. Dudley, S. P. Platania, G.C. White. 2020. Colorado Pikeminnow and Razorback Sucker larval fish survey in the San Juan River during 2020. Final report submitted to San Juan River Implementation Recovery Program, U.S. Fish and Wildlife Service, Albuquerque, NM.
- Fielding, A.H., and J.F. Bell. 1997. A review of methods for assessment of prediction errors in conservation presence/absence models. *Environmental Conservation* 24:38-49.
- Fletcher, D., D. MacKenzie, and E. Villouta. 2005. Modelling skewed data with many zeros: a simple approach combining ordinary and logistic regression. *Environmental and Ecological Statistics* 12: 45-54.
- Franssen, N.R., E.I. Gilbert, and D.L. Propst. 2015. Effects of longitudinal and lateral stream channel complexity on native and non-native fishes in an invaded desert stream. *Freshwater Biology* 60:16-30.
- Gido, K.B., and D.L. Propst. 2012. Long-term dynamics of native and nonnative fishes in the San Juan River, New Mexico and Utah, under a partially managed flow regime. *Transactions of the American Fisheries Society* 141:645-659.
- Peterson, A. T., M. Papes, and J. Soberon. 2008. Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological Modelling* 213:63-72.
- SJRIP (San Juan River Basin Recovery Implementation Program). 2012. San Juan River Basin Recovery Implementation Program monitoring plan and protocols. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- SJRIP (San Juan River Basin Recovery Implementation Program). 2016. Long-range plan. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- USFWS (U.S. Fish and Wildlife Service). 1987. Interagency Standardized Monitoring Program Handbook. Grand Junction, CO.
- Vasconcelos, R. P., O. Le Pape, M. J. Costa, H. N. Cabral. 2013. Predicting estuarine use patterns of juvenile fish with generalized linear models. *Estuarine, Coastal, and Shelf Science* 120:64-74.
- Zeigler, M. P. and M. E. Ruhl. 2016. Small-bodied Fishes Monitoring in the San Juan River: 2015. Final Report submitted to the San Juan River Basin Recovery Implementation Program. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.
- Zeigler, M. P., and M. E. Ruhl. 2017. Small-bodied fishes monitoring in the San Juan River: 2016. Final Report submitted to the San Juan River Basin Recovery Implementation Program. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.

Zeigler, M. P., and J.M. Wick. 2019. Small-bodied fishes monitoring in the San Juan River: 2018. Final Report submitted to the San Juan River Basin Recovery Implementation Program. San Juan River Basin Recovery Implementation Program, U.S. Fish and Wildlife Service, Albuquerque, New Mexico

FY 2022 Budget

Sampling - Animas River to Shiprock, NM

Personnel

Tasks - Annual monitoring of small-bodied fishes in the San Juan River from the Animas River confluence to Shiprock, NM; 1 day trip preparation (8 hrs day) and 4 field days (12 hrs day) = 56 hours (40 hrs regular and 16 hrs overtime).

Project Leader (1)	
40 hrs regular @ \$47.73/hr (\$34.84/hr (base salary) + \$12.89/hr (benefits))	\$ 1,909
16 hrs overtime @ \$71.60/hr (\$47.73/hr * 1.5 (time-and-a-half))	\$ 1,146
Project Biologists (3)	
40 hrs regular @ \$38.24/hr (\$27.91/hr (base salary) + \$10.33 (benefits)) * 3	\$ 4,589
16 hrs overtime @ \$57.36/hr (\$38.24/hr * 1.5 (time-and-a-half)) * 3	\$ 2,753
	<hr/>
Sub-total	\$ 10,397

Per Diem

4 days @ \$151/day (standard NM rate) * 4 biologists	\$ 2,416
	<hr/>
Sub-total	\$ 2,416

Vehicles

Round-trip to Shiprock, NM – 500 miles @ \$0.55/mile	\$ 275
	<hr/>
Sub-total	\$ 275

Animas River to Shiprock Sampling Sub-total \$ 13,088

Sampling - Shiprock, NM to Clay Hills Crossing, UT

Personnel

Tasks - Annual monitoring of small-bodied fishes in the San Juan River from Shiprock, NM to Mexican Hat, UT; The Nature Conservancy RERI Phase I and Phase II sites; 3 days trip preparation (8 hrs day) and 15 field days (12 hrs day) = 204 hours (144 hrs regular and 60 hrs overtime).

Project Leader (1)	
144 hrs regular @ \$47.73/hr (\$34.84/hr (base salary) + \$12.89/hr (benefits))	\$ 6,873
60 hrs overtime @ \$71.60/hr (\$47.73/hr * 1.5 (time-and-a-half))	\$ 4,296
Project Biologist (3)	
144 hrs regular @ \$38.24/hr (\$27.91/hr (base salary) + \$10.33 (benefits)) * 3	\$ 16,520
60 hrs overtime @ \$57.36/hr (\$38.24/hr * 1.5 (time-and-a-half)) * 3	\$ 10,325
	<hr/>
Sub-total	\$ 38,014

Per Diem

15 days @ \$151/day (standard NM rate) * 4 biologists \$ 9,060

Sub-total \$ 9,060

Vehicles

Round-trip (2 vehicles) to Clay Hill Crossing, UT – 1,500 miles @ \$0.55/mile \$ 825

Sub-total \$ 825

Shiprock to Clay Hills Crossing Sampling Sub-total \$ 47,899

Field Equipment & Supplies

Annual equipment and supplies examples: \$ 1,800

- Water quality instrument maintenance 1 @ \$400 \$ 800
- Life Jackets 5 @ \$60 \$ 300
- Raft maintenance \$ 500
- Whirlpacks (500) @ \$50.00/per 500 \$ 50
- Formalin (6 gal) @ \$25/gal \$ 150
- 2 ml cyrogenic vials (1000) @ \$0.40/per 1000 \$ 400
- Ethanol 200 proof (2 gal) @ \$100/gal \$ 200
- Raft (16'), Frame, and Oars (funds not requested) \$ 0

Sub-total \$ 1,800

Sampling Sub-total \$ 62,786

Specimen Management

Personnel

Tasks - Processing (sorting, identification, and data-entry); 12 days of in the laboratory (8 hrs day) = 96 hours.

Project Biologist (1)
96 hrs regular @ \$38.24/hr (\$27.91/hr (base salary) + \$10.33 (benefits)) \$ 3,671

Specimen Management Sub-total \$ 3,671

Data Management/Analysis and Report Preparation

Personnel

Tasks – Data management and QA/QC, data analysis and synthesis, table and graph preparation, report drafting and revision; Project Leader (120 hrs) and one Project Biologist (200 hrs each).

Project Leader (1)
120 hrs regular @ \$47.73/hr (\$34.84/hr (base salary) + \$12.89/hr (benefits)) \$ 5,728

Project Biologist (1)
200 hrs regular @ \$38.24/hr (\$27.91/hr (base salary) + \$10.33 (benefits)) \$ 7,648

Data Management/Analysis & Report Preparation Sub-total \$ 13,376

FY 2022 Total

Sampling Sub-total	\$ 62,786
Specimen Management Sub-total	\$ 3,671
Data Management/Analysis & Report Preparation Sub-total	\$ 13,376
	<hr/>
	Project Sub-Total \$ 79,833
IDC at 22.09 %	\$ 17,635
	<hr/>
	Project Total \$ 97,468